





A
RED
BENCH
IN
PARADISE


RAINBOW
PENSE






TIME






THE COUNTERFEITER WILL BE
PUNISHED WITH PENAL SERVITUDE

GUY BILSON — ALBION — J. H. BILSON



WOW





encore



SANK



WARBLE





de



ar



dde



taedtr69


taed



taed

taed

bliss




indirectly pinned down

A B C E



A QUA S



POESIA, SOLO
POESIA...


the butterfly

ITE

QUCI

qua qua

qua qua




...e tanto amore

oh
oh



o.k.o.k.o.k.o.k.o.k.o.k.o.k.

H de dclins



ISSIMOPOMME

WINE



KNEEW too!
too!
too!
too!




IM
is not
nice
say
H



Y U M



CLARO



13=1°|14=3°|15=5°|16=1°|17=2°|18=3°|19=1°|20=8°|21=2°|22=6°|23=4°|24=2°

I mille segreti

EASE


1=3°|2=5°|3=4°|4=4°|5=2°|6=2°|7=5°|8=2°|9=5°|10=2°|11=4°|12=3°



facile?




DID






FLY



penultimately

S LANDFALL



the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer 1996). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer 1996). The prevalence of schizophrenia in the United States is estimated to be 1.1% (Meltzer 1996).

There is a growing awareness of the need to improve the lives of people with schizophrenia. The World Health Organization (WHO) has developed a set of guidelines for the management of schizophrenia (WHO 1993). The guidelines recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated by a multidisciplinary team, including doctors, nurses, social workers, and psychologists.

DAVID COLE to BETTY DANON
BETTY DANON to DAVID COLE